
PHYSICS

9702/41

Paper 4 A Level Structured Questions

October/November 2016

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2016 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9702	41

- 1 (a) gravitational force provides/is the centripetal force B1
- $$GMm/r^2 = mv^2/r \quad \text{or} \quad GMm/r^2 = mr\omega^2$$
- $$\text{and } v = 2\pi r/T \quad \text{or} \quad \omega = 2\pi/T$$
- M1
- with algebra to $T^2 = 4\pi^2 r^3 / GM$ A1 [3]
- or
- acceleration due to gravity is the centripetal acceleration (B1)
- $$GM/r^2 = v^2/r \quad \text{or} \quad GM/r^2 = r\omega^2$$
- $$\text{and } v = 2\pi r/T \quad \text{or} \quad \omega = 2\pi/T$$
- (M1)
- with algebra to $T^2 = 4\pi^2 r^3 / GM$ (A1)
- (b) (i) equatorial orbit/orbits (directly) above the equator B1
- from west to east B1 [2]
- (ii) $(24 \times 3600)^2 = 4\pi^2 r^3 / (6.67 \times 10^{-11} \times 6.0 \times 10^{24})$ C1
- $$r^3 = 7.57 \times 10^{22}$$
- $$r = 4.2 \times 10^7 \text{ m}$$
- A1 [2]
- (c) $(T/24)^2 = \{(2.64 \times 10^7) / (4.23 \times 10^7)\}^3$ B1
- $$= 0.243$$
- $T = 12$ hours A1 [2]
- or
- $$k (= T^2/r^3) = 24^2 / (4.23 \times 10^7)^3$$
- $$= 7.61 \times 10^{-21}$$
- (B1)
- $$T^2 (= kr^3) = 7.61 \times 10^{-21} \times (2.64 \times 10^7)^3$$
- $$= 140$$
- $T = 12$ hours (A1)
- 2 (a) (i) $p \propto T$ or $pV/T = \text{constant}$ or $pV = nRT$ C1
- $T (= 5 \times 300 =) 1500 \text{ K}$ A1 [2]
- (ii) $pV = nRT$
- $$1.0 \times 10^5 \times 4.0 \times 10^{-4} = n \times 8.31 \times 300$$
- or
- $$5.0 \times 10^5 \times 4.0 \times 10^{-4} = n \times 8.31 \times 1500$$
- C1
- $n = 0.016 \text{ mol}$ A1 [2]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9702	41

- (b) (i) 1. heating/thermal energy supplied B1
2. work done on/to system B1 [2]
- (ii) 1. 240 J A1
2. same value as given in 1. (= 240 J) **and** zero given for 3. A1
3. zero A1 [3]
- 3 (a) $2k/m = \omega^2$ M1
 $\omega = 2\pi f$ M1
 $(2 \times 64 / 0.810) = (2\pi \times f)^2$ leading to $f = 2.0$ Hz A1 [3]
- (b) $v_0 = \omega x_0$ or $v_0 = 2\pi f x_0$
or
 $v = \omega(x_0^2 - x^2)^{1/2}$ and $x = 0$ C1
 $v_0 = 2\pi \times 2.0 \times 1.6 \times 10^{-2}$
 $= 0.20 \text{ ms}^{-1}$ A1 [2]
- (c) frequency: reduced/decreased B1
maximum speed: reduced/decreased B1 [2]
- 4 (a) (i) noise/distortion is removed (from the signal) B1
the (original) signal is reformed/reproduced/recovered/restored B1 [2]
or
signal detected above/below a threshold creates new signal (B1)
of 1s and 0s (B1)
- (ii) noise is superposed on the (displacement of the) signal/cannot be distinguished
or
analogue/signal is continuous (so cannot be regenerated)
or
analogue/signal is not discrete (so cannot be regenerated) B1
noise is amplified with the signal B1 [2]

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9702	41

(b) (i) $\text{gain/dB} = 10 \lg(P_2/P_1)$

$$32 = 10 \lg[P_{\text{MIN}}/(0.38 \times 10^{-6})]$$

or

$$-32 = 10 \lg(0.38 \times 10^{-6}/P_{\text{MIN}})$$

C1

$$P_{\text{MIN}} = 6.0 \times 10^{-4} \text{ W}$$

A1 [2]

(ii) $\text{attenuation} = 10 \lg[(9.5 \times 10^{-3})/(6.02 \times 10^{-4})]$

C1

$$= 12 \text{ dB}$$

$$\text{attenuation per unit length} (= 12/58) = 0.21 \text{ dB km}^{-1}$$

A1 [2]

5 (a) in an electric field, charges (in a conductor) would move

B1

no movement of charge so zero field strength

or

charge moves until $F = 0 / E = 0$

B1 [2]

or

charges in metal do not move

(B1)

no (resultant) force on charges so no (electric) field

(B1)

(b) at P, $E_A = (3.0 \times 10^{-12})/[4\pi\epsilon_0(5.0 \times 10^{-2})^2]$ ($= 10.79 \text{ NC}^{-1}$)

M1

at P, $E_B = (12 \times 10^{-12})/[4\pi\epsilon_0(10 \times 10^{-2})^2]$ ($= 10.79 \text{ NC}^{-1}$)

M1

or

$$(3.0 \times 10^{-12})/[4\pi\epsilon_0(5.0 \times 10^{-2})^2] - (12 \times 10^{-12})/[4\pi\epsilon_0(10 \times 10^{-2})^2] = 0$$

or

$$(3.0 \times 10^{-12})/[4\pi\epsilon_0(5.0 \times 10^{-2})^2] = (12 \times 10^{-12})/[4\pi\epsilon_0(10 \times 10^{-2})^2]$$

(M2)

fields due to charged spheres are (equal and) opposite in direction, so $E = 0$

A1 [3]

(c) $\text{potential} = 8.99 \times 10^9 \{(3.0 \times 10^{-12})/(5.0 \times 10^{-2}) + (12 \times 10^{-12})/(10 \times 10^{-2})\}$

C1

$$= 1.62 \text{ V}$$

A1 [2]

(d) $\frac{1}{2}mv^2 = qV$

$$E_K = \frac{1}{2} \times 107 \times 1.66 \times 10^{-27} \times v^2$$

C1

$$qV = 47 \times 1.60 \times 10^{-19} \times 1.62$$

C1

$$v^2 = 1.37 \times 10^8$$

$$v = 1.2 \times 10^4 \text{ ms}^{-1}$$

A1 [3]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9702	41

- 6 (a) reference to input (voltage) and output (voltage) B1
there is no time delay between change in input and change in output B1 [2]
- or*
- reference to rate at which output voltage changes (B1)
infinite rate of change (of output voltage) (B1)
- (b) (i) $2.00/3.00 = 1.50/R$ C1
- or*
- $V_+ = (3.00 \times 4.5)/(2.00 + 3.00) = 2.7$
 $2.7 = 4.5 \times R/(R + 1.50)$ (C1)
- resistance = 2.25 k Ω A1 [2]
- (ii) 1. correct symbol for LED M1
two LEDs connected with opposite polarities between V_{OUT} and earth A1 [2]
2. below 24 °C, $R_T > 1.5 \text{ k}\Omega$ or resistance of thermistor increases/high B1
- $V_- < V_+$ or V_- decreases/low (must not contradict initial statement) M1
- V_{OUT} is positive/+5 (V) and LED labelled as 'pointing' from V_{OUT} to earth A1 [3]
- 7 (a) region (of space) where a force is experienced by a particle B1 [1]
- (b) (i) gravitational B1
- (ii) gravitational and electric B1
- (iii) gravitational, electric and magnetic B1 [3]
- (c) (i) force (always) normal to direction of motion M1
- (magnitude of) force constant
or
speed is constant/kinetic energy is constant M1
- magnetic force provides/is the centripetal force A1 [3]
- (ii) $mv^2/r = Bqv$ B1
- momentum or p or $mv = Bqr$ B1 [2]

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9702	41

- 8 strong uniform magnetic field B1
- nuclei precess/rotate about field (direction) (1)
- radio-frequency pulse (applied) B1
- R.F. or pulse is at Larmor frequency/frequency of precession (1)
- causes resonance/excitation (of nuclei)/nuclei absorb energy B1
- on relaxation/de-excitation, nuclei emit r.f./pulse B1
- (emitted) r.f./pulse detected and processed (1)
- non-uniform magnetic field B1
- allows position of nuclei to be located B1
- allows for location of detection to be changed/different slices to be studied (1)
- any two of the points marked (1)* B2 [8]
- 9 (a) (induced) e.m.f. proportional to rate of change of (magnetic) flux (linkage) M1
A1 [2]
- (b) flux linkage = BAN
- $$= \pi \times 10^{-3} \times 2.8 \times \pi \times (1.6 \times 10^{-2})^2 \times 85 = 6.0 \times 10^{-4} \text{ Wb}$$
- B1 [1]
- (c) e.m.f. = $\Delta N\Phi / \Delta t$
- $$= (6.0 \times 10^{-4} \times 2) / 0.30$$
- C1
- $$= 4.0 \text{ mV}$$
- A1 [2]
- (d) sketch: $E = 0$ for $t = 0 \rightarrow 0.3 \text{ s}$, $0.6 \text{ s} \rightarrow 1.0 \text{ s}$, $1.6 \text{ s} \rightarrow 2.0 \text{ s}$ B1
- $E = 4 \text{ mV}$ for $t = 0.3 \text{ s} \rightarrow 0.6 \text{ s}$ (either polarity) B1
- $E = 2 \text{ mV}$ for $t = 1.0 \text{ s} \rightarrow 1.6 \text{ s}$ B1
- with opposite polarity B1 [4]

Page 7	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9702	41

- 10 (a)** electromagnetic radiation/photons incident on a surface B1
causes emission of electrons (from the surface) B1 [2]
- (b)** $E = hc / \lambda$
 $= (6.63 \times 10^{-34} \times 3.00 \times 10^8) / (436 \times 10^{-9})$ C1
 $= 4.56 \times 10^{-19} \text{ J } (4.6 \times 10^{-19} \text{ J})$ A1 [2]
- (c) (i)** $\Phi = hc / \lambda_0$
 $\lambda_0 = (6.63 \times 10^{-34} \times 3.00 \times 10^8) / (1.4 \times 1.60 \times 10^{-19})$ C1
 $= 890 \text{ nm}$ A1 [2]
- (ii)** $\lambda_0 = (6.63 \times 10^{-34} \times 3.00 \times 10^8) / (4.5 \times 1.60 \times 10^{-19})$
 $= 280 \text{ nm}$ A1 [1]
- (d)** caesium:
wavelength of photon less than threshold wavelength (or v.v.)
or
 $\lambda_0 = 890 \text{ nm} > 436 \text{ nm}$
so yes A1
- tungsten:
wavelength of photon greater than threshold wavelength (or v.v.)
or
 $\lambda_0 = 280 \text{ nm} < 436 \text{ nm}$
so no A1 [2]
- 11** in metal, conduction band overlaps valence band/no forbidden band/no band gap B1
as temperature rises, no increase in number of free electrons/charge carriers B1
as temperature rises, lattice vibrations increase M1
(lattice) vibrations restrict movement of electrons/charge carriers M1
(current decreases) so resistance increases A1 [5]

Page 8	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – October/November 2016	9702	41

- 12 (a) (i) time for number of atoms/nuclei or activity to be reduced to one half M1
- reference to (number of...) original nuclide/single isotope
or
reference to half of original value/initial activity A1 [2]
- (ii) $A = A_0 \exp(-\lambda t)$ and either $t = t_{1/2}$, $A = \frac{1}{2}A_0$ or $\frac{1}{2}A_0 = A_0 \exp(-\lambda t_{1/2})$ M1
- so $\ln 2 = \lambda t_{1/2}$ (and $\ln 2 = 0.693$), hence $0.693 = \lambda t_{1/2}$ A1 [2]
- (b) $A = \lambda N$
- $N = 200 / (2.1 \times 10^{-6})$ C1
- $= 9.52 \times 10^7$ C1
- mass = $(9.52 \times 10^7 \times 222 \times 10^{-3}) / (6.02 \times 10^{23})$
- or*
- mass = $9.52 \times 10^7 \times 222 \times 1.66 \times 10^{-27}$ C1
- $= 3.5 \times 10^{-17} \text{ kg}$ A1 [4]